

$\Lambda(1520)$ D_{03} $I(J^P) = 0(\frac{3}{2}^-)$ Status: ***

Discovered by FERRO-LUZZI 62; the elaboration in WATSON 63 is the classic paper on the Breit-Wigner analysis of a multichannel resonance.

The measurements of the mass, width, and elasticity published before 1975 are now obsolete and have been omitted. They were last listed in our 1982 edition Physics Letters **111B** 1 (1982).

Production and formation experiments agree quite well, so they are listed together here.

 $\Lambda(1520)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1519.5 ±1.0 OUR ESTIMATE				
1519.50±0.18 OUR AVERAGE				
1517.3 ±1.5	300	BARBER	80D	SPEC $\gamma p \rightarrow \Lambda(1520) K^+$
1519 ±1		GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
1517.8 ±1.2	5k	BARLAG	79	HBC $K^- p$ 4.2 GeV/c
1520.0 ±0.5		ALSTON-...	78	DPWA $\bar{K}N \rightarrow \bar{K}N$
1519.7 ±0.3	4k	CAMERON	77	HBC $K^- p$ 0.96–1.36 GeV/c
1519 ±1		GOPAL	77	DPWA $\bar{K}N$ multichannel
1519.4 ±0.3	2000	CORDEN	75	DBC $K^- d$ 1.4–1.8 GeV/c

 $\Lambda(1520)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
15.6 ±1.0 OUR ESTIMATE				
15.59±0.27 OUR AVERAGE				
16.3 ±3.3	300	BARBER	80D	SPEC $\gamma p \rightarrow \Lambda(1520) K^+$
16 ±1		GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
14 ±3	677	¹ BARLAG	79	HBC $K^- p$ 4.2 GeV/c
15.4 ±0.5		ALSTON-...	78	DPWA $\bar{K}N \rightarrow \bar{K}N$
16.3 ±0.5	4k	CAMERON	77	HBC $K^- p$ 0.96–1.36 GeV/c
15.0 ±0.5		GOPAL	77	DPWA $\bar{K}N$ multichannel
15.5 ±1.6	2000	CORDEN	75	DBC $K^- d$ 1.4–1.8 GeV/c

 $\Lambda(1520)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 N\bar{K}$	45 ± 1%
$\Gamma_2 \Sigma\pi$	42 ± 1%
$\Gamma_3 \Lambda\pi\pi$	10 ± 1%
$\Gamma_4 \Sigma(1385)\pi$	

Γ_5	$\Sigma(1385)\pi(\rightarrow \Lambda\pi\pi)$
Γ_6	$\Lambda(\pi\pi)_S$ -wave
Γ_7	$\Sigma\pi\pi$
Γ_8	$\Lambda\gamma$
Γ_9	$\Sigma^0\gamma$

CONSTRAINED FIT INFORMATION

An overall fit to 9 branching ratios uses 26 measurements and one constraint to determine 6 parameters. The overall fit has a $\chi^2 = 17.6$ for 21 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-64				
x_3	-32 -34				
x_7	-4	-3	-1		
x_8	-8	-7	-3	0	
x_9	-24	-21	-10	-1	-1
	x_1	x_2	x_3	x_7	x_8

$\Lambda(1520)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.


$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.45 ± 0.01 OUR ESTIMATE			
0.447 ± 0.007 OUR FIT	Error includes scale factor of 1.2.		
0.455 ± 0.011 OUR AVERAGE			
0.47 ± 0.02	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
0.45 ± 0.03	ALSTON-...	78	DPWA $\bar{K}N \rightarrow \bar{K}N$
0.448 ± 0.014	CORDEN	75	DBC $K^- d$ 1.4–1.8 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.47 ± 0.01	GOPAL	77	DPWA See GOPAL 80
0.42	MAST	76	HBC $K^- p \rightarrow \bar{K}^0 n$

$\Gamma(\Sigma\pi)/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.42 ± 0.01 OUR ESTIMATE			
0.420 ± 0.007 OUR FIT	Error includes scale factor of 1.2.		
0.423 ± 0.011 OUR AVERAGE			
0.426 ± 0.014	CORDEN	75	DBC $K^- d$ 1.4–1.8 GeV/c
0.418 ± 0.017	BARBARO-...	69B	HBC $K^- p$ 0.28–0.45 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.46	KIM	71	DPWA K-matrix analysis

$\Gamma(\Sigma\pi)/\Gamma(N\bar{K})$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
0.940±0.026 OUR FIT	Error includes scale factor of 1.3.			
0.95 ±0.04 OUR AVERAGE	Error includes scale factor of 1.7. See the ideogram below.			
0.98 ± 0.03	² GOPAL 77	DPWA	$\bar{K}N$ multichannel	
0.82 ± 0.08	BURKHARDT 69	HBC	$K^- p$ 0.8–1.2 GeV/c	
1.06 ± 0.14	SCHEUER 68	DBC	$K^- N$ 3 GeV/c	
0.96 ± 0.20	DAHL 67	HBC	$\pi^- p$ 1.6–4 GeV/c	
0.73 ± 0.11	DAUBER 67	HBC	$K^- p$ 2 GeV/c	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.06 ± 0.12	BERTHON 74	HBC	Quasi-2-body σ	
1.72 ± 0.78	MUSGRAVE 65	HBC		

WEIGHTED AVERAGE
0.95±0.04 (Error scaled by 1.7)



$\Gamma(\Lambda\pi\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
0.10 ±0.01 OUR ESTIMATE				
0.095±0.005 OUR FIT	Error includes scale factor of 1.2.			
0.096±0.008 OUR AVERAGE	Error includes scale factor of 1.6.			
0.091±0.006	CORDEN 75	DBC	$K^- d$ 1.4–1.8 GeV/c	
0.11 ± 0.01	³ MAST 73B	IPWA	$K^- p \rightarrow \Lambda\pi\pi$	

$\Gamma(\Lambda\pi\pi)/\Gamma(N\bar{K})$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
0.213±0.012 OUR FIT	Error includes scale factor of 1.2.			
0.202±0.021 OUR AVERAGE				
0.22 ± 0.03	BURKHARDT 69	HBC	$K^- p$ 0.8–1.2 GeV/c	
0.19 ± 0.04	SCHEUER 68	DBC	$K^- N$ 3 GeV/c	
0.17 ± 0.05	DAHL 67	HBC	$\pi^- p$ 1.6–4 GeV/c	
0.21 ± 0.18	DAUBER 67	HBC	$K^- p$ 2 GeV/c	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.27 ± 0.13	BERTHON 74	HBC	Quasi-2-body σ	
0.2	KIM 71	DPWA	K-matrix analysis	

$\Gamma(\Sigma\pi)/\Gamma(\Lambda\pi\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_3
4.42±0.25 OUR FIT	Error includes scale factor of 1.2.			
3.9 ± 0.6 OUR AVERAGE				
3.9 ± 1.0	UHLIG 67	HBC	$K^- p$ 0.9–1.0 GeV/c	
3.3 ± 1.1	BIRMINGHAM 66	HBC	$K^- p$ 3.5 GeV/c	
4.5 ± 1.0	ARMENTEROS65C	HBC		

$\Gamma(\Sigma(1385)\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
0.041±0.005	CHAN 72	HBC	$K^- p \rightarrow \Lambda\pi\pi$	

$\Gamma(\Sigma(1385)\pi(\rightarrow \Lambda\pi\pi))/\Gamma(\Lambda\pi\pi)$

Γ_5/Γ_3

The $\Lambda\pi\pi$ mode is largely due to $\Sigma(1385)\pi$. Only the values of $(\Sigma(1385)\pi)/(\Lambda\pi\pi)$ given by MAST 73B and CORDEN 75 are based on real 3-body partial-wave analyses. The discrepancy between the two results is essentially due to the different hypotheses made concerning the shape of the $(\pi\pi)_S$ -wave state.

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_3
0.58±0.22	CORDEN 75	DBC	$K^- d$ 1.4–1.8 GeV/c	
0.82±0.10	⁴ MAST 73B	IPWA	$K^- p \rightarrow \Lambda\pi\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.39±0.10	⁵ BURKHARDT 71	HBC	$K^- p \rightarrow (\Lambda\pi\pi)\pi$	

$\Gamma(\Lambda(\pi\pi)_S\text{-wave})/\Gamma(\Lambda\pi\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_3
0.20±0.08	CORDEN 75	DBC	$K^- d$ 1.4–1.8 GeV/c	

$\Gamma(\Sigma\pi\pi)/\Gamma_{\text{total}}$

Γ_7/Γ

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
0.009 ± 0.001 OUR ESTIMATE				
0.0086±0.0005 OUR FIT				
0.0086±0.0005 OUR AVERAGE				
0.007 ± 0.002	⁶ CORDEN 75	DBC	$K^- d$ 1.4–1.8 GeV/c	
0.0085±0.0006	⁷ MAST 73	MPWA	$K^- p \rightarrow \Sigma\pi\pi$	
0.010 ± 0.0015	BARBARO-... 69B	HBC	$K^- p$ 0.28–0.45 GeV/c	

$\Gamma(\Lambda\gamma)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.5 ± 1.5 OUR ESTIMATE				
8.8 ± 1.1 OUR FIT				
8.8 ± 1.1 OUR AVERAGE				
10.7 \pm 2.9 $^{+1.5}_{-0.4}$	32	TAYLOR 05	CLAS	$\gamma p \rightarrow K^+ \Lambda\gamma$
10.2 \pm 2.1 \pm 1.5	290	ANTIPOV 04A	SPNX	$p N(C) \rightarrow \Lambda(1520) K^+ N(C)$
8.0 \pm 1.4	238	MAST 68B	HBC	Using $\Gamma(N\bar{K})/\Gamma_{\text{total}} = 0.45$

 $\Gamma(\Sigma^0\gamma)/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0195 ± 0.0034 OUR FIT			
0.02 \pm 0.0035	⁸ MAST	68B HBC	Not measured; see note

 $\Lambda(1520)$ FOOTNOTES1 From the best-resolution sample of $\Lambda\pi\pi$ events only.2 The $\bar{K}N \rightarrow \Sigma\pi$ amplitude at resonance is $+0.46 \pm 0.01$.3 Assumes $\Gamma(N\bar{K})/\Gamma_{\text{total}} = 0.46 \pm 0.02$.4 Both $\Sigma(1385)\pi DS_{03}$ and $\Sigma(\pi\pi) DP_{03}$ contribute.5 The central bin (1514–1524 MeV) gives 0.74 ± 0.10 ; other bins are lower by 2-to-5 standard deviations.6 Much of the $\Sigma\pi\pi$ decay proceeds via $\Sigma(1385)\pi$.7 Assumes $\Gamma(N\bar{K})/\Gamma_{\text{total}} = 0.46$.8 Calculated from $\Gamma(\Lambda\gamma)/\Gamma_{\text{total}}$, assuming SU(3). Needed to constrain the sum of all the branching ratios to be unity. **$\Lambda(1520)$ REFERENCES**

TAYLOR	05	PR C71 054609	S. Taylor <i>et al.</i>	(JLab CLAS Collab.)
Also		PR C72 039902 (Errat.)	S. Taylor <i>et al.</i>	(JLab CLAS Collab.)
ANTIPOV	04A	PL B604 22	Yu.M. Antipov <i>et al.</i>	(IHEP SPHINX Collab.)
PDG	82	PL 11B 1	M. Roos <i>et al.</i>	(HELS, CIT, CERN)
BARBER	80D	ZPHY C7 17	D.P. Barber <i>et al.</i>	(DARE, LANC, SHEF)
GOPAL	80	Toronto Conf. 159	G.P. Gopal	(RHEL) IJP
BARLAG	79	NP B149 220	S.J.M. Barlag <i>et al.</i>	(AMST, CERN, NIJM+)
ALSTON-...	78	PR D18 182	M. Alston-Garnjost <i>et al.</i>	(LBL, MTJO+) IJP
Also		PRL 38 1007	M. Alston-Garnjost <i>et al.</i>	(LBL, MTJO+) IJP
CAMERON	77	NP B131 399	W. Cameron <i>et al.</i>	(RHEL, LOIC) IJP
GOPAL	77	NP B119 362	G.P. Gopal <i>et al.</i>	(LOIC, RHEL) IJP
MAST	76	PR D14 13	T.S. Mast <i>et al.</i>	(LBL)
CORDEN	75	NP B84 306	M.J. Corden <i>et al.</i>	(BIRM)
BERTHON	74	NC 21A 146	A. Berthon <i>et al.</i>	(CDEF, RHEL, SACL+)
MAST	73	PR D7 3212	T.S. Mast <i>et al.</i>	(LBL) IJP
MAST	73B	PR D7 5	T.S. Mast <i>et al.</i>	(LBL) IJP
CHAN	72	PRL 28 256	S.B. Chan <i>et al.</i>	(MASA, YALE)
BURKHARDT	71	NP B27 64	E. Burkhardt <i>et al.</i>	(HEID, CERN, SACL)
KIM	71	PRL 27 356	J.K. Kim	(HARV) IJP
Also		Duke Conf. 161	J.K. Kim	(HARV) IJP
Hyperon Resonances, 1970				
BARBARO-...	69B	Lund Conf. 352	A. Barbaro-Galtieri <i>et al.</i>	(LRL)
Also		Duke Conf. 95	R.D. Tripp	(LRL)
Hyperon Resonances 1970				
BURKHARDT	69	NP B14 106	E. Burkhardt <i>et al.</i>	(HEID, EFI, CERN+)
MAST	68B	PRL 21 1715	T.S. Mast <i>et al.</i>	(LRL)

SCHEUER	68	NP B8 503	J.C. Scheuer <i>et al.</i>	(SABRE Collab.)
DAHL	67	PR 163 1377	O.I. Dahl <i>et al.</i>	(LRL)
DAUBER	67	PL 24B 525	P.M. Dauber <i>et al.</i>	(UCLA)
UHLIG	67	PR 155 1448	R.P. Uhlig <i>et al.</i>	(UMD, NRL)
BIRMINGHAM	66	PR 152 1148	M. Haque <i>et al.</i>	(BIRM, GLAS, LOIC, OXF+)
ARMENTEROS	65C	PL 19 338	R. Armenteros <i>et al.</i>	(CERN, HEID, SACL)
MUSGRAVE	65	NC 35 735	B. Musgrave <i>et al.</i>	(BIRM, CERN, EPOL+)
WATSON	63	PR 131 2248	M.B. Watson, M. Ferro-Luzzi, R.D. Tripp	(LRL) IJP
FERRO-LUZZI	62	PRL 8 28	M. Ferro-Luzzi, R.D. Tripp, M.B. Watson	(LRL) IJP
